

ALMOST SURE INVARIANCE PRINCIPLE FOR NON-AUTONOMOUS HOLOMORPHIC DYNAMICS IN $\mathbb{P}^k$

TURGAY BAYRAKTAR

ABSTRACT. We prove almost sure invariance principle, a strong form of approximation by Brownian motion, for non-autonomous holomorphic dynamical systems on complex projective space $\mathbb{P}^k$ for Hölder continuous and DSH observables.

1. INTRODUCTION

Let $f : \mathbb{P}^k \rightarrow \mathbb{P}^k$ be a holomorphic map of algebraic degree $d \geq 2$ and ω_{FS} denote the Fubini-Study form on $\mathbb{P}^k$ normalized by $\int \omega_{FS}^k = 1$. Dynamical Green current T_f of f is defined to be the weak limit of the sequence of smooth forms $\{d^{-n}(f^n)^*\omega_{FS}\}$ ([Bro65, HP94, FS95]). Green currents play an important role in the dynamical study of holomorphic endomorphisms of the projective space [FS95, Sib99]. The current T_f has Hölder continuous quasi-potentials, hence by Bedford-Taylor theory the exterior products

$$T_f^p = T_f \wedge \cdots \wedge T_f$$

are also well-defined for $1 \leq p \leq k$ and dynamically interesting currents. In particular, the top degree intersection $\mu_f = T_f^k$ yields the unique f -invariant measure of maximal entropy ([Lju83, Sib99, BD01]) with many interesting stochastic properties. For instance, in [Dup10] Dupont obtained an almost sure principle (ASIP) for the holomorphic dynamical system $(\mathbb{P}^k, \mathcal{B}, f, \mu_f)$ by using coding techniques and applying Philipp-Stout's theorem [PS75] for observables with analytic singularities. The coding techniques were originally introduced by Przytycki-Urbański-Zdunik [PUZ89] in complex dimension one from which they deduced ASIP (see also [Hay99, PRL07] and the references therein for some statistical results in the case of dimension one). Recall that ASIP in the context of a holomorphic dynamical system indicates that for suitable class of observables ψ , the partial Birkhoff sums $\frac{1}{n} \sum_{j=0}^{n-1} \psi \circ f^j$ can be approximated by a Brownian motion (at integer times) in such a way that almost surely the error between the trajectories is negligible relative to the size of the trajectories (see §3 for details). Some important statistical laws such as Central Limit Theorem (CLT) and Law of Iterated Logarithm (LIL) are among the immediate consequences of ASIP. In this context, CLT was also obtained by [CLB05, DS06a, DNS10] by means of different methods. The approach of Dinh-Nguyen-Sibony [DNS10] is based on Gordin's martingale approximation method [Gor69] and exponential decay of correlations for the Hölder continuous and DSH observables. Recall that a DSH function can be locally written as a difference of two plurisubharmonic (psh) functions.

In [Bay15], we studied ergodic and statistical properties of random dynamical systems of holomorphic endomorphisms of $\mathbb{P}^k$. The latter is defined as successive iterations of holomorphic maps which are chosen at random according to a fixed probability measure. Under a mild assumption on the probability measure, there is a naturally associated ergodic skew-product and a stationary

2000 *Mathematics Subject Classification.* 37F10, 60F17, 32H50.

Key words and phrases. Almost Sure Invariance Principle, Central Limit Theorem, Holomorphic maps, Non-autonomous dynamical systems.

homogenous Markov chain in that dynamical setting for which we established two versions of CLT. The first one is for the partial Birkhoff sums of the skew product for Hölder continuous and DSH observables. The second one is for the backward images of randomly chosen points with respect to the stationary probability law of the chain.

In the present paper, we consider non-autonomous holomorphic dynamical systems of the complex projective space $\mathbb{P}^k$. The latter is defined as successive iterations of a sequence of non-linear holomorphic maps of the same algebraic degree. In this setting, we obtain more refined statistical results. Namely, we prove an ASIP (Theorem 1.2) for these non-stationary systems under a natural assumption (1.1) on the distance of the tail of the sequence from the complement of holomorphic maps. It should be emphasized that the martingale approximation method of Gordin [Gor69] gives rise to a reverse martingale increment sequence. Then using the martingale CLT, one can deduce CLT for the observations of a stationary dynamical system. The ASIP can often also be obtained in this way for the class of systems which are closed under time-reversal (see [MN05] and references therein). However, for classes of non-stationary systems that are intrinsically time-orientated, such as non-autonomous holomorphic dynamics considered in in this paper, the situation can be more delicate (see eg. [MN05, CM15, HNTV17] for examples of real dynamical systems). Here, we utilize exponential decay of correlations for DSH and Hölder continuous observables and the abstract invariance principle obtained by [CM15] in order to prove that non-autonomous holomorphic dynamical systems satisfy ASIP.

1.1. Main Result. Recall that the set of holomorphic endomorphisms $\mathcal{H}_d$ of fixed algebraic degree d is a Zariski open subset of the set of meromorphic maps $\mathcal{M}_d$ and the complement $\mathcal{M} := \mathcal{M}_d \setminus \mathcal{H}_d$ is an irreducible hypersurface [GKZ94]. The set $\mathcal{M}_d$ can be identified with $\mathbb{P}^N$ where $N = (k+1)\binom{d+k}{d} - 1$. In what follows, we consider $\mathbb{P}^N$ as a metric space furnished with the Fubini-Study metric $\text{dist}(\cdot, \cdot)$.

For a sequence $\mathbf{f} := (f_0, f_1, \dots)$ of holomorphic maps $f_n : \mathbb{P}^k \rightarrow \mathbb{P}^k$ of fixed algebraic degree $d \geq 2$ we study statistical properties of the iterates

$$F_n := f_{n-1} \circ \dots \circ f_1 \circ f_0 : \mathbb{P}^k \rightarrow \mathbb{P}^k.$$

In [Bay15] (see also [DT13]), for such a sequence $\mathbf{f} = (f_j) \subset \mathcal{H}_d$ whose “tail” is sufficiently far from the complement $\mathcal{M}$, we proved the existence of a measure $\mu_{\mathbf{f}}$ which describes asymptotic distribution of pre-images of a generic point $z \in \mathbb{P}^k$.

Theorem 1.1. [Bay15] *Let $\mathbf{f} = (f_j)_{j \geq 0} \subset \mathcal{H}_d$ be a sequence of holomorphic maps such that*

$$(1.1) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \log \text{dist}(f_j, \mathcal{M}) = C > -\infty.$$

Then there exists a Borel measure $\mu_{\mathbf{f}}$ such that for every smooth measure ν on $\mathbb{P}^k$

$$F_n^* \nu \rightarrow \mu_{\mathbf{f}}$$

in the sense of measures as $n \rightarrow \infty$. The measure $\mu_{\mathbf{f}}$ has Hölder continuous potentials.

In §2, we provide an alternative proof for Hölder continuity of local potentials of invariant measures $\mu_{\mathbf{f}}$ given by Theorem 1.1. Furthermore, in Theorem 2.7 we obtain strong mixing properties of sequential holomorphic dynamical system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$. As a consequence, we prove that the system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$ is exact, a strong form of ergodicity (Theorem 2.5).

Our main result gives an Almost Sure Invariance Principle (ASIP) for sequences of holomorphic endomorphisms verifying hypothesis (1.1) and for sequences of dsh and Hölder continuous observables.

Theorem 1.2 (ASIP). *Let $\mathbf{f} = (f_0, f_1, \dots)$ be a sequence of holomorphic maps of fixed algebraic degree $d \geq 2$ verifying (1.1). Let also ψ_n be a sequence of DSH (respectively, α -Hölder continuous) functions satisfying $\sup_n \|\psi_n\|_{DSH} < \infty$ (respectively, $\sup_n \|\psi_n\|_{\mathcal{C}_\alpha} < \infty$). Assume that $\int_{\mathbb{P}^k} \psi_n d\mu_n = 0$ and the variances $\sigma_n^2 := \int_{\mathbb{P}^k} (\sum_{j=0}^{n-1} \psi_j \circ F_j)^2 d\mu_{\mathbf{f}}$ satisfy*

$$(1.2) \quad n = O(\sigma_n^2).$$

Then on an extended probability space there exists a sequence of centered independent Gaussian random variables $(Z_j)_{j \in \mathbb{N}}$ such that $\mathbb{E}[Z_j^2] = \sigma_n^2(1 + o(1))$ and that

$$(1.3) \quad \left| \sum_{j=0}^{n-1} \psi_j \circ F_j - \sum_{j=0}^{n-1} Z_j \right| = O(n^{1-\gamma}) \quad \mu_{\mathbf{f}} \text{ a.e.}$$

for some $0 < \gamma < \frac{1}{2}$.

Some remarks in order. The assumption $\int_{\mathbb{P}^k} \psi_n d\mu_n = 0$ is not restrictive as we may replace ψ_n with $\psi_n - \int_{\mathbb{P}^k} \psi_n d\mu_n$. An immediate consequence of Theorem 1.2 is CLT that is

$$(1.4) \quad \frac{1}{\sigma_n} \sum_{j=0}^{n-1} \psi_j \circ F_j \rightarrow \mathcal{N}(0, 1)$$

in distribution $\mu_{\mathbf{f}}$ (cf [PS75]). Another consequence is LIL which implies that

$$\limsup_n \left[\frac{1}{\sqrt{\sigma_n \log \log(\sigma_n)}} \sum_{j=0}^{n-1} \psi_j \circ F_j \right] = 1 \quad \mu_{\mathbf{f}} - a.e.$$

and

$$\liminf_n \left[\frac{1}{\sqrt{\sigma_n \log \log(\sigma_n)}} \sum_{j=0}^{n-1} \psi_j \circ F_j \right] = -1 \quad \mu_{\mathbf{f}} - a.e.$$

Finally, we remark that for an autonomous holomorphic dynamical system (i.e. $f = f_j \forall j$) and a single observable (i.e. $\psi = \psi_j, \forall j$) we have either σ_n is bounded (in this case ψ is a coboundary i.e. $\psi = \zeta - \zeta \circ f$ for some $\zeta \in L_{\mu_f}^2$) or the variances σ_n is of order $\sqrt{n}$. Hence, we recover [Dup10, Theorem C].

2. ERGODICITY AND MIXING FOR NON-AUTONOMOUS SYSTEMS

2.1. Invariant measures. For a sequence of endomorphisms $\mathbf{f} = (f_j)_{j \geq 0} \subset \mathcal{H}_d$ we define *topological Lyapunov exponent*

$$\chi_{top}(\mathbf{f}) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \sup_{x \in \mathbb{P}^k} \|D_x F_j\|$$

where $\|\cdot\|$ is a fixed norm. Clearly, this definition does not depend on the choice of the norm. Note that the limit exists (possibly infinite) due to the sub-multiplicity of the sequence $\sup_{x \in \mathbb{P}^k} \|D_x F_j\|$. The next lemma will be useful in the sequel:

Lemma 2.1. *Let $\mathbf{f} = (f_j)_{j \geq 0} \subset \mathcal{H}_d$ verifying (1.1). Then $0 < \chi_{top}(\mathbf{f}) < \infty$.*

Proof. Note that

$$\Psi : \mathcal{M}_d \rightarrow \mathcal{M}_d$$

$$\Psi(g, x) = g(x)$$

defines a meromorphic map. Then it follows from [DD04, Lemma 2.1] that there exists $C > 0$ and $q \geq 1$ such that

$$\|D_x f_j\| \leq \|D_{f_j, x} \Psi\| \leq C \text{dist}(f_j, \mathcal{M})^{-q}, \quad \forall j \geq 0.$$

Then

$$(2.1) \quad M_n := \sup_{x \in \mathbb{P}^k} \|D_x F_n\| \leq C^n \prod_{j=0}^{n-1} \text{dist}(f_j, \mathcal{M})^{-q}$$

and this in turn implies that

$$\frac{1}{n} \log \left(\sup_{x \in \mathbb{P}^k} \|D_x F_n\| \right) \leq C - \frac{q}{n} \sum_{j=0}^{n-1} \log \text{dist}(f_j, \mathcal{M}).$$

□

The following result was motivated by [Sib99, Remark 1.7.2].

Theorem 2.2. *Let $\mathbf{f} = (f_j)_{j \geq 0} \subset \mathcal{H}_d$ verifying (1.1). Then*

$$\frac{1}{n} F_n^* \omega_{FS} \rightarrow \omega_{FS} + dd^c g_{\mathbf{f}}$$

in the sense of currents. Moreover, Green function $g_{\mathbf{f}}$ is α -Hölder continuous for every exponent $\alpha < \frac{\log d}{\chi_{\text{top}}(\mathbf{f})}$.

Proof. First, we sketch the proof of existence of the limit. Let u_j be a smooth qpsH function defined by

$$\frac{1}{d} f_j^* \omega_{FS} = \omega_{FS} + dd^c u_j.$$

Then

$$\begin{aligned} \frac{1}{d^n} F_n^* \omega_{FS} &= \frac{1}{d^n} f_0^* \cdots f_{n-1}^* \omega_{FS} \\ &= \omega_{FS} + dd^c \sum_{j=0}^{n-1} \frac{1}{d^j} u_j \circ F_j \\ &=: \omega_{FS} + dd^c g_n \end{aligned}$$

By [DT13, Proposition 3] we have

$$(2.2) \quad \|u_j\|_{C^1} \leq C \text{dist}(f_j, \mathcal{M})^{-q}.$$

It follows from (1.1) that for each $\epsilon > 0$ there exists $j_0 \in \mathbb{N}$ such that

$$(2.3) \quad \text{dist}(f_j, \mathcal{M}) \geq e^{-\epsilon j} \quad \forall j \geq j_0.$$

Hence,

$$\begin{aligned} \sup_{x \in \mathbb{P}^k} |g_n(x) - g_m(x)| &= \left| \sum_{j=n}^{m-1} \frac{1}{d^j} u_j \circ F_j \right| \\ &\leq \sum_{j=n}^{m-1} \frac{\sup_{x \in \mathbb{P}^k} |u_j(x)|}{d^j} \\ &\leq C \sum_{j=n}^{m-1} \frac{e^{\epsilon q j}}{d^j}. \end{aligned}$$

This implies that $g_n \rightarrow g_{\mathbf{f}}$ uniformly on $\mathbb{P}^k$ for some continuous qpsH function $g_{\mathbf{f}}$.

In order to prove Hölder continuity, let $0 < \alpha < \frac{\log d}{\chi_{\text{top}}(\mathbf{f})}$. Note that by (2.1)

$$\text{dist}(F_n(x), F_n(y)) \leq C^n M_n \text{dist}(x, y).$$

Then by (2.2) and (2.3) for small $\epsilon > 0$ we obtain

$$\begin{aligned} |g_{\mathbf{f}}(x) - g_{\mathbf{f}}(y)| &\leq \sum_{j \geq 0} \frac{|u_j(F_j(x)) - u_j(F_j(y))|}{d^j} \\ &\leq \sum_{j \geq 0} \frac{C \text{dist}(f_j, \mathcal{M})^{-q} \text{dist}(F_j(x), F_j(y))^\alpha}{d^j} \\ &\leq C \text{dist}(x, y)^\alpha \sum_{j \geq 0} \frac{\text{dist}(f_j, \mathcal{M})^{-q} C^{\alpha j} e^{\chi_{\text{top}}(\mathbf{f}) \alpha j}}{d^j} \\ &\leq C_1 \text{dist}(x, y)^\alpha. \end{aligned}$$

□

Since the *dynamical Green current* $T_{\mathbf{f}}$ given by Theorem 2.2 has Hölder continuous potentials by Bedford-Taylor theory the exterior powers

$$T_{\mathbf{f}}^p := T_{\mathbf{f}} \wedge \cdots \wedge T_{\mathbf{f}}$$

are also well-defined positive closed (p, p) currents for each $1 \leq p \leq k$. Letting T_n^p to be the dynamical Green current associated with $\mathbf{f}_n := (f_n, f_{n+1}, \dots)$ it is easy to see that they inherit the invariance properties

$$(2.4) \quad f_{n-1}^* T_n^p = d^p T_n^p \text{ and } (f_{n-1})_* T_{n-1}^p = d^{k-p} T_n^p \quad \forall n \geq 1$$

in particular,

$$(2.5) \quad f_{n-1}^* \mu_n = d^k \mu_{n-1} \text{ and } (f_{n-1})_* \mu_{n-1} = \mu_n \quad \forall n \geq 1.$$

The next result is a direct consequence of [DT13, Proposition 11] and Theorem 1.1:

Proposition 2.3. *The top degree dynamical Green current*

$$T_{\mathbf{f}}^k = T_{\mathbf{f}} \wedge \cdots \wedge T_{\mathbf{f}}$$

coincides with the invariant measure $\mu_{\mathbf{f}}$. In particular, $\mu_{\mathbf{f}}$ has continuous Hölder potentials.

2.2. Sequential Ergodicity, Mixing and Exactness. In what follows we let $\mathcal{B}$ denote the Borel algebra on $\mathbb{P}^k$ and $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$ be a *sequential holomorphic dynamical system* that is $\mathbf{f} = (f_0, f_1, \dots)$ is a sequence of holomorphic maps $f_j \in \mathcal{H}_d$ and we assume that (1.1) holds. We say that the system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$ is *mean ergodic* if

$$(2.6) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} [\mu_{\mathbf{f}}(B \cap F_j^{-1}(A)) - \mu_{\mathbf{f}}(B) \mu_j(A)] = 0$$

for all $\mathcal{B}$ -measurable sets A, B . Note that (2.6) is equivalent to

$$(2.7) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} \left[\int_{\mathbb{P}^k} \varphi \cdot \psi \circ F_j d\mu_{\mathbf{f}} - \int_{\mathbb{P}^k} \varphi d\mu_{\mathbf{f}} \int_{\mathbb{P}^k} \psi d\mu_j \right] = 0$$

for all continuous (equivalently smooth, bounded or $L_{\mu_{\mathbf{f}}}^2$) functions φ, ψ on $\mathbb{P}^k$. Using the property $P_j 1 = 1$ and the argument in [CM15, Remark 1.3] we see that (2.6) is equivalent to convergence in L^q -norm for $1 \leq q < \infty$:

$$(2.8) \quad \lim_{n \rightarrow \infty} \left\| \frac{1}{n} \sum_{j=0}^{n-1} \psi \circ F_j - \int \psi d\mu_{\mathbf{f}} \right\|_{L^q_{\mu_{\mathbf{f}}}} \rightarrow 0.$$

Finally, we remark that in the setting of autonomous dynamical systems the mean ergodicity is equivalent to Birkhoff's ergodic theorem to which we refer here as point-wise ergodic. However, for non-autonomous systems these two concepts are not equivalent (cf. [CM15]). We say that the sequential holomorphic dynamical system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$ is *point-wise ergodic* if

$$(2.9) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} [\psi \circ F_j(x) - \int \psi d\mu_{\mathbf{f}}] = 0 \quad \mu_{\mathbf{f}} - a.s.$$

for all $L^2_{\mu_{\mathbf{f}}}$ functions ψ on $\mathbb{P}^k$.

We say that the system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$ is *mixing* if for all smooth functions φ, ψ on $\mathbb{P}^k$ we have

$$(2.10) \quad \int_{\mathbb{P}^k} \varphi \cdot \psi \circ F_n d\mu_{\mathbf{f}} - \int \varphi d\mu_{\mathbf{f}} \int \psi d\mu_{\mathbf{f}} \rightarrow 0.$$

Clearly, sequential mixing implies mean ergodicity (2.7). In Theorem 2.7, we prove that every holomorphic sequential dynamical system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$ is mixing and hence mean ergodic. In fact, we obtain a stronger form of mean ergodicity.

For a given sequence $\mathbf{f} = (f_j)_{j \geq 0}$ of holomorphic maps $f_j \in \mathcal{H}_d$ we define

$$(2.11) \quad \mathcal{B}_j := F_j^{-1}(\mathcal{B}) = f_0^{-1} \cdots f_{j-1}^{-1}(\mathcal{B}).$$

Note that

$$\mathcal{B} = \mathcal{B}_0 \supset \mathcal{B}_1 \supset \cdots \supset \mathcal{B}_n.$$

We also denote the asymptotic σ -algebra

$$\mathcal{B}_{\infty} := \bigcap_{j=0}^{\infty} \mathcal{B}_j.$$

Definition 2.4. A sequential holomorphic dynamical system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$ satisfying (1.1) is called *exact* if the asymptotic σ -algebra

$$\mathcal{B}_{\infty} = \{\emptyset, \mathbb{P}^k\}$$

modulo $\mu_{\mathbf{f}}$.

It follows from the definition of exact sequence that $\mathbf{f}$ is exact if and only if

$$\bigcap_{j=0}^{\infty} L^2_{\mu_{\mathbf{f}}}(\mathcal{B}_j) = \{\text{constant functions}\}$$

where

$$L^2_{\mu_{\mathbf{f}}}(\mathcal{B}_j) = \{\xi \in L^2_{\mu_{\mathbf{f}}}(\mathbb{P}^k) : \xi \text{ is } \mathcal{B}_j \text{ measurable}\}.$$

Clearly, exactness implies mixing hence, mean ergodicity.

In what follows we let $\mathbf{f} = (f_j)_{j \geq 0} \subset \mathcal{H}_d$ satisfying (1.1). Then each $f_j \in \mathcal{H}_d$ induces a unitary operator

$$\begin{aligned} T_j : L^2_{\mu_{j+1}}(\mathbb{P}^k) &\rightarrow L^2_{\mu_j}(\mathbb{P}^k) \\ T_j(\varphi) &= \varphi \circ f_j. \end{aligned}$$

We denote the adjoint of this operator by

$$\begin{aligned} P_j : L^2_{\mu_j}(\mathbb{P}^k) &\rightarrow L^2_{\mu_{j+1}}(\mathbb{P}^k) \\ P_j \psi(x) &= d^{-k} \sum_{f_j(y)=x} \psi(y) \end{aligned}$$

and let

$$\mathcal{P}_j = P_{j-1} \cdots P_1 P_0.$$

Theorem 2.5. *Let $\mathbf{f} = (f_j)_{j \geq 0}$ be a sequence of holomorphic maps in $\mathcal{H}_d$ satisfying (1.1). Then the system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$ is exact.*

We remark that Theorem 2.5 generalizes [Pet05] in which exactness was obtained for compact for sequences $\mathbf{f} = (f_j)$ i.e.

$$\text{dist}(f_j, \mathcal{M}) \geq C > 0, \forall j \geq 0.$$

The proof given in [Pet05] is based on showing that for each F_j there are sufficiently many inverse branches on a disc away from the critical values for which the pre-images has small diameter. The latter is originally due to Briend and Duval [BD01]. This method breaks down in the present setting and we provide a different approach.

Proof. Note that another equivalent condition for exactness of the sequence $\mathbf{f} = (f_j)_{j \geq 0}$ is

$$(2.12) \quad \|\mathcal{P}_j \psi\|_{L^1_{\mu_j}} \rightarrow 0 \text{ as } j \rightarrow \infty$$

for every $\psi \in L^1_{\mu_{\mathbf{f}}}$ such that $\|\psi\|_{L^1_{\mu_{\mathbf{f}}}} = 0$. Indeed, given such ψ it follows from Doob's martingale convergence theorem that the conditional expectations

$$\mathbb{E}[\psi | \mathcal{B}_j] \rightarrow \mathbb{E}[\psi | \mathcal{B}_{\infty}] \text{ in } L^1_{\mu_{\mathbf{f}}}$$

as $j \rightarrow \infty$. Since $\mathbb{E}[\psi | \mathcal{B}_j] = (\mathcal{P}_j \psi) \circ F_j$ by invariance properties (2.5) we have

$$\|\mathbb{E}[\psi | \mathcal{B}_j]\|_{L^1_{\mu_{\mathbf{f}}}} = \|\mathcal{P}_j \psi\|_{L^1_{\mu_j}}$$

and the claim follows.

Finally, as $P_j 1 \equiv 1 \forall j$, it is enough to verify the condition (2.12) for smooth functions and this follows from Lemma 2.10 below. $\square$

Sequential Mixing. In this section we explore mixing properties of the sequential holomorphic dynamical system $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_{\mathbf{f}})$. We start with some preliminaries:

2.2.1. DSH Functions. Recall that a function φ is call a *quassi-plurisubharmonic* (qpsh for short) if φ can be locally written as sum of a smooth function and a pluri-subharmonic (psh) function. We say that a function $\psi \in L^1(\mathbb{P}^k)$ is dsh if outside a pluripolar set $\psi = \varphi_1 - \varphi_2$ where φ_i are qpsh functions. This implies that

$$dd^c \psi = T^+ - T^-$$

for some positive closed $(1, 1)$ currents $T^{\pm}$. Two dsh functions are identified if they coincide outside a pluripolar set; we denote the set of all dsh functions by $DSH(\mathbb{P}^k)$. Note that dsh functions are stable under pull-back and push-forward operators induced by meromorphic self-maps of $\mathbb{P}^k$ and have good compactness properties inherited from those of qpsh functions. Following [DS06b] one can define a norm on $DSH(\mathbb{P}^k)$ as follows:

$$\|\psi\|_{DSH} := \|\psi\|_{L^1(\mathbb{P}^k)} + \inf \|T^{\pm}\|$$

where $dd^c \psi = T^+ - T^-$ and the infimum is taken over all such representations. We remark that the currents $T^{\pm}$ have the same mass as they are cohomologous and $\|\cdot\|_{DSH} \lesssim \|\cdot\|_{\mathcal{C}^2}$. Moreover, it follows from properties of psh functions that $\|\cdot\|_{L^p} \lesssim \|\cdot\|_{DSH}$ for $1 \leq p < \infty$.

If μ is a probability measure on $\mathbb{P}^k$ such that all qpsh functions are μ -integrable then one can define

$$\|\psi\|_{DSH}^{\mu} := |\langle \mu, \psi \rangle| + \inf \|T^{\pm}\|$$

where $T^{\pm}$ as above and $\langle \mu, \psi \rangle := \int_{\mathbb{P}^k} \psi d\mu$.

Proposition 2.6. [DS06b] *Let $\psi \in DSH(\mathbb{P}^k)$ then there exists negative qps functions φ_1, φ_2 such that $\psi = \varphi_1 - \varphi_2$ and $dd^c \varphi_i \geq -c\|\psi\|_{DSH}\omega$ where $c > 0$ independent of ψ and φ_i 's. Moreover, $|\psi|$ is also a dsh function and $\|\psi\|_{DSH} \leq c\|\psi\|_{DSH}$.*

Now, we turn our attention to strong mixing properties of sequential holomorphic dynamical systems.

Theorem 2.7. *Let $(\mathbb{P}^k, \mathcal{B}, \mathbf{f}, \mu_f)$ be a sequential holomorphic dynamical system satisfying (1.1). Then for every $\varphi \in L^p_{\mu_n}(\mathbb{P}^k)$ and $\psi \in DSH(\mathbb{P}^k)$ we have*

$$|\langle \mu_f, (\varphi \circ f_{n-1} \circ \cdots \circ f_0)\psi \rangle - \langle \mu_n, \varphi \rangle \langle \mu_f, \psi \rangle| \leq Cd^{-n} \|\varphi\|_{L^p_{\mu_n}} \|\psi\|_{DSH}^{\mu_f}$$

where $C > 0$ depends only on $\mathbf{f}$ and $p > 1$. Moreover, for each $0 \leq \alpha \leq 2$ there exists $C > 0$ depending only on $\mathbf{f}$ such that

$$|\langle \mu_f, (\varphi \circ f_{n-1} \circ \cdots \circ f_0)\psi \rangle - \langle \mu_n, \varphi \rangle \langle \mu_f, \psi \rangle| \leq Cd^{-\frac{\alpha n}{2}} \|\varphi\|_{L^p_{\mu_n}} \|\psi\|_{\mathcal{C}^\alpha}$$

for each $\varphi \in L^p_{\mu_n}$ and ψ of class $\mathcal{C}^\alpha$.

We need several preliminary lemmas to prove Theorem 2.7. In what follows, C_f denotes a constant which depends only on $\mathbf{f}$.

Lemma 2.8. *There exists $C_f > 0$ such that*

$$\|\psi\|_{DSH}^{\mu_n} \leq C_f \|\psi\|_{DSH}$$

for every $n \in \mathbb{N}$ and $\psi \in DSH(\mathbb{P}^k)$.

Proof. Let $\psi \in DSH(\mathbb{P}^k)$ then by Proposition 2.6 there exists qps functions ϕ_i such that $\psi = \phi_1 - \phi_2$ and $dd^c \phi_i \geq -c\|\psi\|_{DSH}\omega_{FS}$ where $c > 0$ is independent of ψ and ϕ_i . Since the measures μ_n have Hölder continuous super potentials (cf. [Bay15, Theorem 1.1]) with Hölder exponent $0 < \alpha \leq 1$, by [DN14, Lemma 3.3] we obtain

$$|\langle \mu_n, \psi \rangle| \leq C_f \max(\|\psi\|_{L^1}, c^{1-\alpha} \|\psi\|_{DSH}^{1-\alpha} \|\psi\|_{L^1}^\alpha)$$

If $\|\psi\|_{L^1} \geq c^{1-\alpha} \|\psi\|_{DSH}^{1-\alpha} \|\psi\|_{L^1}^\alpha$ we are done. Otherwise $\|\psi\|_{L^1} < c\|\psi\|_{DSH}$ and this implies that

$$|\langle \mu_n, \psi \rangle| \leq cC_f \|\psi\|_{DSH}.$$

Thus, the assertion follows from Proposition 2.6. $\square$

Remark 2.9. *By using a similar argument and using Lemma 2.8 one can also show that there exists a constant $C_f > 0$ such that*

$$\|\psi\|_{DSH} \leq C_f \|\psi\|_{DSH}^{\mu_n}.$$

for every $n \geq 0$ (cf. [DT13, Proposition 8]).

The following lemma is essentially due to [DNS10], however, we need to make some modifications to adapt it in our setting.

Lemma 2.10. *There exists $C_f > 0$ such that for every $\psi \in DSH(\mathbb{P}^k)$ satisfying*

$$\langle \mu_f, \psi \rangle = 0$$

and for every $q \geq 1$

$$(2.13) \quad \|\mathcal{P}_n(\psi)\|_{L^q_{\mu_j}} \leq qC_f d^{-n} \|\psi\|_{DSH}^{\mu_f}$$

for all $j, n \geq 1$.

Proof. Note that $(f_{n-1})^* \mu_n = d^k \mu_{n-1}$ for $n \geq 1$. This implies that

$$\langle \mu_n, P_{n-1}(P_{n-2} \circ \cdots \circ P_0 \psi) \rangle = 0.$$

Moreover,

$$\|P_{n-1}(P_{n-2} \circ \cdots \circ P_0 \psi)\|_{DSH}^{\mu_n} \leq d^{-1} \|P_{n-2} \circ \cdots \circ P_0 \psi\|_{DSH}^{\mu_{n-1}}$$

Indeed, we may write

$$dd^c(P_{n-2} \circ \cdots \circ P_0 \psi) = R_{n-2}^+ - R_{n-2}^-$$

where $R_{n-2}^\pm$ are some positive closed $(1, 1)$ currents. Then

$$(2.14) \quad \|P_{n-1}(P_{n-2} \circ \cdots \circ P_0 \psi)\|_{DSH}^{\mu_n} \leq \|d^{-k}(f_{n-1})_*(R_{n-2}^\pm)\| = d^{-1} \|R_{n-2}^\pm\|$$

where the last equality follows from cohomological computation. Now by Proposition 2.6, Remark 2.9, Lemma 2.8 and (2.14) we obtain

$$\begin{aligned} \|P_{n-1} \circ \cdots \circ P_1 \circ P_0(\psi)\|_{DSH} &\leq C \|P_{n-1} \circ \cdots \circ P_1 \circ P_0(\psi)\|_{DSH} \\ &\leq C_1 \|P_{n-1} \circ \cdots \circ P_1 \circ P_0(\psi)\|_{DSH}^{\mu_n} \\ &\leq C_2 d^{-n} \|\psi\|_{DSH}^{\mu_f} \\ &\leq C_3 d^{-n} \|\psi\|_{DSH} \end{aligned}$$

where $C_3 > 0$ depends on $\mathbf{f}$ but does not depend on n nor ψ . Thus, by above estimate and Lemma 2.8 it is enough to prove the case $\|\psi\|_{DSH} > 0$. Since

$$\frac{d^n}{\|\psi\|_{DSH}} |P_{n-1} \circ \cdots \circ P_1 \circ P_0(\psi)|$$

is a bounded sequence in $DSH(\mathbb{P}^k)$, by Theorem 1.1 and [DNS10, Corollary 1.2] (see also [DN14, Proposition 4.4]) there exists $\gamma > 0$ and $C_f > 0$ independent of ψ such that

$$\langle \mu_j, \exp(\gamma \frac{d^n}{\|\psi\|_{DSH}} |P_{n-1} \circ \cdots \circ P_1 \circ P_0(\psi)|) \rangle \leq C_f$$

for all $j, n \geq 1$. Finally, by using the inequality $\frac{x^q}{q!} \leq e^x$ for $x \geq 0$ we conclude that

$$\|P_{n-1} \circ \cdots \circ P_1 \circ P_0(\psi)\|_{L_{\mu_j}^q} \leq q C_{f,\gamma} d^{-n} \|\psi\|_{DSH}.$$

□

In the deterministic case, as a consequence of interpolation theory between the Banach spaces $\mathcal{C}^0$ and $\mathcal{C}^2$ [Tri78]; it was observed in [DNS10] that a holomorphic map $f \in \mathcal{H}_d$ posses strong mixing property for α -Hölder continuous functions with $0 < \alpha \leq 1$ (see [DNS10, Proposition 3.5]). Adapting their argument to our setting, we obtain the succeeding lemma. We omit the proof as it is similar to the one given in Lemma 2.10 and to that of [DNS10, Proposition 3.5].

Lemma 2.11. *Let $1 \leq q < \infty$ and $0 < \alpha \leq 1$ be fixed. If $\psi : \mathbb{P}^k \rightarrow \mathbb{R}$ be a α -Hölder continuous function satisfying $\langle \mu_f, \psi \rangle = 0$ then there exists a constant $C_{f,\alpha} > 0$ independent of ψ such that*

$$\|\mathcal{P}_n(\psi)\|_{L^q(\mu_n)} \leq C_{f,\alpha} q^{\frac{\alpha}{2}} d^{-\frac{n\alpha}{2}} \|\psi\|_{\mathcal{C}^\alpha}$$

for every $n \geq 1$.

Proof of Theorem 2.7. If ψ is constant then the assertion follows from the invariance properties

$$(f_j)_* \mu_j = \mu_{j+1}.$$

Thus, replacing ψ by $\psi - \langle \mu_{\mathbf{f}}, \psi \rangle$ we may assume that $\langle \mu_{\mathbf{f}}, \psi \rangle = 0$. Then by Hölder's inequality and applying Lemma 2.10 with $q = \frac{p}{p-1}$ we obtain

$$\begin{aligned} |\langle \mu_{\mathbf{f}}, (\varphi \circ f_{n-1} \circ \cdots \circ f_0) \psi \rangle| &= d^{-kn} |\langle F_n^* \mu_n, (\varphi \circ f_{n-1} \circ \cdots \circ f_0) \psi \rangle| \\ &\leq |\langle \mu_n, \varphi P_{n-1} \circ \cdots \circ P_1 \circ P_0(\psi) \rangle| \\ &\leq \|\varphi\|_{L^p(\mu_n)} \|P_{n-1} \circ \cdots \circ P_1 \circ P_0(\psi)\|_{L^q(\mu_n)} \\ &\leq \frac{p}{p-1} C_{\mathbf{f}} d^{-n} \|\varphi\|_{L^p(\mu_n)} \|\psi\|_{DSH}^{\mu_{\mathbf{f}}} \end{aligned}$$

for some $c > 0$ independent of ψ and for all $n \geq 0$.

Finally, repeating the above argument by using Lemma 2.11 the second assertion follows. $\square$

The following result follows from Theorem 2.7; its proof is based on induction and Hölder's inequality. As the proof is similar to that of [DNS10, Theorem 3.4] we omit it.

Corollary 2.12. *Let $\mathbf{f} = (f_j)_{j \in \mathbb{N}}$ and μ_j be as in Theorem 2.7 and $r \geq 1$ be an integer. Further, we let ψ_j be dsh functions satisfying $\sup_{0 \leq j \leq r} \|\psi_j\|_{DSH} < \infty$. Then*

$$|\langle \mu_{\mathbf{f}}, \psi_0(\psi_1 \circ F_{n_1}) \cdots (\psi_r \circ F_{n_r}) \rangle - \prod_{j=0}^r \langle \mu_{n_j}, \psi_j \rangle| \leq C_{\mathbf{f}} d^{-n} \prod_{j=0}^r \|\psi_j\|_{DSH}$$

where $0 = n_0 \leq n_1 \leq \cdots \leq n_r$ and $n := \min_{0 \leq j \leq r} (n_{j+1} - n_j)$.

Note that we may also obtain an analogue statement for Hölder continuous functions. Next, we obtain a strong law of large numbers (SLLN) for dsh and Hölder continuous observables. This result will be used to establish ASIP (Theorem 1.2).

Theorem 2.13. *Let $\{\phi_n\}_{n \in \mathbb{N}}$ be a sequence of dsh (respectively, Hölder continuous functions with exponent $0 < \alpha \leq 1$) such that $\sup_{n \in \mathbb{N}} \|\phi_n\|_{DSH} < \infty$ (respectively, $\sup_{n \in \mathbb{N}} \|\phi_n\|_{\mathcal{C}^\alpha} < \infty$) and $\langle \mu_n, \phi_n \rangle = 0$ for $n \geq 0$. Then for each integer $r \geq 1$ we have*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} (\phi_j^r \circ F_j - \langle \phi_j^r, \mu_j \rangle) = 0 \text{ } \mu_{\mathbf{f}}\text{-a.s.}$$

Proof. We prove the theorem for dsh functions as Hölder continuous case is similar. Letting

$$X_j := \phi_j^r \circ F_j - \langle \mu_j, \phi_j^r \rangle \text{ for } j \geq 0$$

by (2.5) we see that $\mathbb{E}[X_j] := \int_{\mathbb{P}^k} X_j d\mu_{\mathbf{f}} = 0$. Note that by Lemma 2.8 we have $\|\phi_j\|_{L_{\mu_j}^p} \lesssim \|\phi_j\|_{DSH}$ for $p \geq 1$ and we infer that

$$\text{Var}[X_j] := \mathbb{E}[X_j^2] - \mathbb{E}[X_j]^2 = \mathbb{E}[X_j^2] = O(1).$$

Recall that the covariance of X_j and X_ℓ is given by

$$\text{Cov}(X_j, X_\ell) := \mathbb{E}[X_j X_\ell] - \mathbb{E}[X_j] \mathbb{E}[X_\ell] = \mathbb{E}[X_j X_\ell].$$

Moreover, denoting $m_j := \langle \mu_j, \phi_j \rangle$ by (2.5), triangle inequality and Corollary 2.12 we have

$$\begin{aligned} |\text{Cov}(X_j, X_{j+\ell})| &\leq \left| \int_{\mathbb{P}^k} \phi_j^r \circ F_j \cdot \phi_{j+\ell}^r \circ F_{j+\ell} d\mu_{\mathbf{f}} \right| + |m_j m_{j+\ell}| \\ &\lesssim d^{-\ell} \|\phi_j\|_{DSH}^r \|\phi_{j+\ell}\|_{DSH}^r + d^{-2j-\ell} \|\phi_j\|^r \|\phi_{j+\ell}\|^r \\ &\lesssim d^{-\ell} \|\phi_j\|_{DSH}^r \|\phi_{j+\ell}\|_{DSH}^r (1 + d^{-2j}). \end{aligned}$$

where the implied constant does not depend on ℓ .

Hence, the assertion follows from [Lyo88, Corollary 11]. $\square$

3. ASIP

First, we recall some basic notions that we will need in the sequel. Let $(U_j)_{j \in \mathbb{N}}$ be a sequence of random variables on a probability space $(X, \mathcal{F}, \mu)$ such that $\mathbb{E}[U_j] = 0 \ \forall j$. We say that $(U_j)_{j \in \mathbb{N}}$ satisfies almost sure invariance principle (ASIP) if there exists a sequence of independent centered Gaussian random variables $(Z_j)_{j \in \mathbb{N}}$ such that on an extended probability space

$$(3.1) \quad \sum_{j=0}^{n-1} U_j = \sum_{j=0}^{n-1} Z_j + O(\sigma_n^{1-\gamma}) \ \mu - a.e$$

where $\gamma \in (0, 1)$ a fixed constant and $\sigma_n = \sum_{j=0}^{n-1} \mathbb{E}[Z_j^2] \rightarrow \infty$.

Recall that a Brownian motion at integer times coincides with a sum of i.i.d. Gaussian variables, hence the above definition can also be formulated as an almost sure approximation by a Brownian motion, with error $O(\sigma_n^{1-\gamma})$.

A sequence of random variables $(U_j)_{j \in \mathbb{N}}$ is called a *reversed martingale difference* if there exists a non-increasing sequence $(\mathcal{F}_j)_{j \in \mathbb{N}}$ of σ -algebras (i.e. $\mathcal{F}_{j+1} \subset \mathcal{F}_j$) such that

- (1) U_j is $\mathcal{F}_j$ -measurable
- (2) $\mathbb{E}[U_j | \mathcal{F}_{j+1}] = 0 \ \forall j \geq 0$.

The next lemma will be useful in the proof of Theorem 1.2.

Lemma 3.1. [CM15, Lemma 4.2] *Let $(\zeta_n)_{n \in \mathbb{N}}$ be a sequence of reversed martingale differences in L^p_μ , $1 \leq p \leq 2$ with respect to a non-increasing filtration $(\mathcal{F}_j)_{j \in \mathbb{N}}$. Assume that*

$$\sum_{n \geq 0} \mathbb{E}[|\zeta_n|^p] < \infty.$$

Then $\sum_{n \geq 0} \zeta_n$ converges μ -a.s. and in L^p .

We will use the following abstract ASIP for sequences of reversed martingale differences:

Theorem 3.2. [CM15] *Let $(U_j)_{j \in \mathbb{N}}$ be a sequence of square integrable random variables adapted to a non-increasing filtration $(\mathcal{F}_j)$. Assume that $\mathbb{E}[U_j | \mathcal{F}_{j+1}] = 0 \ \mu$ -a.s and $\sigma_n^2 = \sum_{j=0}^{n-1} \mathbb{E}[U_j^2] \rightarrow \infty$ and that $\sup_j \mathbb{E}[U_j^2] < \infty$. Let also a_n be a non-decreasing sequence of positive real numbers such that*

$$(a_n / \sigma_n^2)_{n \in \mathbb{N}} \text{ is non-increasing and } (a_n / \sigma_n)_{n \in \mathbb{N}} \text{ is non-decreasing.}$$

Assume further that

$$(3.2) \quad \sum_{j=0}^{n-1} (\mathbb{E}[U_j^2 | \mathcal{F}_{j+1}] - \mathbb{E}[U_j^2]) = o(a_n) \ \mu - a.s.$$

$$(3.3) \quad \sum_{j=0}^{\infty} a_n^{-q} \mathbb{E}[|U_j|^{2q}] < \infty \text{ for some } 1 \leq q \leq 2.$$

Then on an extended probability space there exists a sequence $(Z_j)_{j \in \mathbb{N}}$ of independent centered Gaussian variables such that $\mathbb{E}[Z_j^2] = \mathbb{E}[U_j^2]$ and

$$(3.4) \quad \sup_{0 \leq k \leq n-1} \left| \sum_{j=0}^k U_j - \sum_{j=0}^k Z_j \right| = o([a_n (|\log(\frac{\sigma_n^2}{a_n})| + \log \log a_n)]^{\frac{1}{2}}) \ \mu - a.s.$$

Proof of Theorem 1.2. We prove the Theorem for DSH observables as Hölder continuous case is the similar. Note that

$$\mathcal{P}_n(\psi_n \circ F_n) = \psi_n.$$

Moreover, since $\mathbb{E}[\phi|\mathcal{B}_n]$ is orthogonal projection of $\phi \in L^2(\mathcal{B}, \mu_{\mathbf{f}})$ to the Hilbert space $L^2(\mathcal{B}_n, \mu_{\mathbf{f}})$ it is easy to see that

$$(3.5) \quad \mathbb{E}[\psi_n \circ F_n | \mathcal{B}_n] = (\mathcal{P}_n \psi_n) \circ F_n \quad \mu_{\mathbf{f}} - a.s.$$

This implies that

$$(3.6) \quad \|\mathbb{E}[\psi_n \circ F_n | \mathcal{B}_n]\|_{L^2_{\mu_{\mathbf{f}}}} = \|\mathcal{P}_n \psi_n\|_{L^2_{\mu_n}} \leq C_{\mathbf{f}} d^{-n} \|\psi_n\|_{DSH}^{\mu_{\mathbf{f}}}$$

where the latter inequality follows from Lemma 2.10.

Now, we define

$$h_j = P_{j-1}(\psi_{j-1} + h_{j-1}) \quad \forall j \geq 1$$

and set $h_0 \equiv 0$. We also let

$$U_j := (\psi_j + h_j - h_{j+1} \circ f_j) \circ F_j \quad \forall j \geq 0.$$

Note that since $P_j(\psi_j + h_j) - h_{j+1} = 0$, by (3.5) we obtain

$$\mathbb{E}[U_j | \mathcal{B}_{j+1}] = 0$$

Hence, $(U_j)_{j \in \mathbb{N}}$ form a sequence of reversed martingale differences for the filtration $(\mathcal{B}_j)_{j \in \mathbb{N}}$ defined by (2.11). Moreover,

$$(3.7) \quad \sum_{j=0}^{n-1} U_j = S_n - h_n \circ F_n$$

where $S_n := \sum_{j=0}^{n-1} \psi_j \circ F_j$. Then by invariance properties (2.5) and Lemma 2.10 we have

$$\|h_n \circ F_n\|_{L^2_{\mu_{\mathbf{f}}}} = \|h_n\|_{L^2_{\mu_n}} \leq C_1$$

where $C_1 > 0$ depends only on $\mathbf{f}$ but independent of n . This implies that

$$\begin{aligned} \left| \|S_n\|_{L^2_{\mu_{\mathbf{f}}}} - \left\| \sum_{j=0}^{n-1} U_j \right\|_{L^2_{\mu_{\mathbf{f}}}} \right| &\leq \left\| S_n - \sum_{j=0}^{n-1} U_j \right\|_{L^2_{\mu_{\mathbf{f}}}} \\ &= \|h_n \circ F_n\|_{L^2_{\mu_{\mathbf{f}}}} \leq C_1 \end{aligned}$$

and we conclude that $\left\| \sum_{j=0}^{n-1} U_j \right\|_{L^2_{\mu_{\mathbf{f}}}}^2 \rightarrow \infty$. Then using $\int_{\mathbb{P}^k} U_i U_j d\mu_{\mathbf{f}} = 0$ for $i \neq j$ we conclude that

$$\sum_{j=0}^{n-1} \mathbb{E}[U_j^2] = \left\| \sum_{j=0}^{n-1} U_j \right\|_{L^2_{\mu_{\mathbf{f}}}}^2 \rightarrow \infty$$

where $\mathbb{E}[U_j^2] = \int_{\mathbb{P}^k} U_j^2 d\mu_{\mathbf{f}}$. We also remark that by Lemma 2.10

$$(3.8) \quad \sup_j \|U_j\|_{L^q_{\mu_{\mathbf{f}}}} \leq C_q, \quad \forall q \geq 1.$$

Hence, it suffices to verify hypothesis (3.2) and (3.3) of Theorem 3.2. To this end we can choose an non-decreasing sequence of positive real numbers $C_1 n \leq a_n \leq C_2 n$ such that (a_n/σ_n^2) is non-increasing and $(a_n/\sigma_n) \uparrow \infty$. Note that

$$(3.9) \quad \sum_{j \geq 0} \frac{\|U_j\|_{L^4_{\mu_{\mathbf{f}}}}^4}{a_j^2} \leq \sum_{j \geq 0} \frac{C_3}{n^2} < \infty$$

which verifies (3.3) with $q = 2$. Moreover, by (3.9) and Lemma 3.1

$$\sum_{j=0}^{\infty} \frac{\mathbb{E}[U_j^2 | \mathcal{B}_{j+1}] - U_j^2}{j}$$

converges $\mu_{\mathbf{f}}$ -a.s. Then by Kronecker's Lemma

$$\sum_{j=0}^{n-1} (\mathbb{E}[U_j^2 | \mathcal{B}_{j+1}] - U_j^2) = o(n) \text{ } \mu_{\mathbf{f}}\text{-a.s.}$$

Hence, in order to get (3.3), it is enough to show that

$$(3.10) \quad \sum_{j=0}^{n-1} (U_j^2 - \mathbb{E}[U_j^2]) = o(n) \text{ } \mu_{\mathbf{f}}\text{-a.s.}$$

Let us denote by

$$\phi_j := (\psi_j + h_j - h_{j+1} \circ f_j)$$

so that $U_j^2 = \phi_j^2 \circ F_j$ for $j \geq 0$. Note that ϕ_j are dsh functions and by Lemma 2.10 their dsh norms are bounded. Moreover, since $\langle \mu_j, h_j \rangle = 0$ by (2.5) we have $\langle \mu_j, \phi_j \rangle = 0$ for $j \geq 0$. Hence, the claim (3.10) follows from Theorem 2.13.

For the Hölder continuous observables, Theorem 2.13 does not apply directly to the functions ϕ_j . In this case, first we expand $\phi_j^2 \circ F_j$ and utilize Lemma 2.11 to estimate the covariances as in Theorem 2.13 and obtain SLLN.

Finally, applying Theorem 3.2 on an extended probability space we obtain centered independent Gaussian random variables such that

$$\sup_{0 \leq k \leq n-1} \left| \sum_{j=0}^k U_j - \sum_{j=0}^k Z_j \right| = o([n(O(1) + \log \log n)]^{\frac{1}{2}}) \text{ } \mu_{\mathbf{f}}\text{-a.s.}$$

□

REFERENCES

- [Bay15] T. Bayraktar, *Ergodic properties of random holomorphic endomorphisms of $\mathbb{P}^k$* , Int. Math. Res. Not. IMRN (2015), no. 4, 927–959.
- [Bay15a] ———, *Corrigendum to: Ergodic properties of random holomorphic endomorphisms of $\mathbb{P}^k$ [MR3340342]*, Int. Math. Res. Not. IMRN (2015), no. 14, 6005–6009. MR 3384466
- [BD01] J.-Y. Briend and J. Duval, *Deux caractérisations de la mesure d'équilibre d'un endomorphisme de $\mathbb{P}^k(\mathbb{C})$* , Publ. Math. Inst. Hautes Études Sci. (2001), no. 93, 145–159. MR 1863737 (2002k:32027)
- [Bro65] H. Brolin, *Invariant sets under iteration of rational functions*, Ark. Mat. **6** (1965), 103–144 (1965). MR 0194595 (33 #2805)
- [CLB05] S. Cantat and S. Le Borgne, *Théorème limite central pour les endomorphismes holomorphes et les correspondances modulaires*, Int. Math. Res. Not. (2005), no. 56, 3479–3510. MR 2200586 (2007c:60024)
- [CM15] C. Cuny and F. Merlevède, *Strong invariance principles with rate for “reverse” martingale differences and applications*, J. Theoret. Probab. **28** (2015), no. 1, 137–183. MR 3320963
- [DD04] T.-C. Dinh and C. Dupont, *Dimension de la mesure d'équilibre d'applications méromorphes*, J. Geom. Anal. **14** (2004), no. 4, 613–627. MR 2111420 (2006k:37117)
- [DN14] T.-C. Dinh and V.-A. Nguyễn, *Characterization of Monge-Ampère measures with Hölder continuous potentials*, J. Funct. Anal. **266** (2014), no. 1, 67–84. MR 3121721
- [DNS10] T.C. Dinh, V.A. Nguyễn, and N. Sibony, *Exponential estimates for plurisubharmonic functions*, Journal of Differential Geometry **84** (2010), no. 3, 465–488.
- [DS06a] T.-C. Dinh and N. Sibony, *Decay of correlations and the central limit theorem for meromorphic maps*, Comm. Pure Appl. Math. **LIX** (2006), no. 0754–0768, 754–768.

- [DS06b] ———, *Distribution des valeurs de transformations méromorphes et applications*, Comment. Math. Helv. **81** (2006), no. 1, 221–258. MR 2208805 (2007i:32017)
- [DT13] H. De Thélin, *Endomorphismes pseudo-aléatoires dans les espaces projectifs. I*, Manuscripta Math. **142** (2013), no. 3-4, 347–367.
- [Dup10] C. Dupont, *Bernoulli coding map and almost sure invariance principle for endomorphisms of $\mathbb{P}^k$* , Probab. Theory Related Fields **146** (2010), no. 3-4, 337–359. MR 2574731 (2010k:37088)
- [FS95] J. E. Fornæss and N. Sibony, *Complex dynamics in higher dimension. II*, Modern methods in complex analysis (Princeton, NJ, 1992), Ann. of Math. Stud., vol. 137, Princeton Univ. Press, Princeton, NJ, 1995, pp. 135–182. MR 1369137 (97g:32033)
- [GKZ94] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky, *Discriminants, resultants, and multidimensional determinants*, Mathematics: Theory & Applications, Birkhäuser Boston Inc., Boston, MA, 1994. MR 1264417 (95e:14045)
- [Gor69] M. I. Gordin, *The central limit theorem for stationary processes*, Dokl. Akad. Nauk SSSR **188** (1969), 739–741. MR 0251785 (40 #5012)
- [Hay99] N. Haydn, *Convergence of the transfer operator for rational maps*, Ergodic Theory Dynam. Systems **19** (1999), no. 3, 657–669. MR 1695914
- [HNTV17] N. Haydn, M. Nicol, A. Török, and S. Vaienti, *Almost sure invariance principle for sequential and non-stationary dynamical systems*, Trans. of the Amer. Math. Soc. (2017).
- [HP94] J. H. Hubbard and P. Papadopol, *Superattractive fixed points in $\mathbb{C}^n$* , Indiana Univ. Math. J. **43** (1994), no. 1, 321–365. MR 1275463 (95e:32025)
- [Lju83] M. Ju. Ljubich, *Entropy properties of rational endomorphisms of the Riemann sphere*, Ergodic Theory Dynam. Systems **3** (1983), no. 3, 351–385. MR 741393 (85k:58049)
- [Lyo88] R. Lyons, *Strong laws of large numbers for weakly correlated random variables*, Michigan Math. J. **35** (1988), no. 3, 353–359. MR 978305
- [MN05] I. Melbourne and M. Nicol, *Almost sure invariance principle for nonuniformly hyperbolic systems*, Comm. Math. Phys. **260** (2005), no. 1, 131–146. MR 2175992
- [Pet05] H. Peters, *Non-autonomous dynamics in $\mathbb{P}^k$* , Ergodic Theory Dynam. Systems **25** (2005), no. 4, 1295–1304. MR 2158406 (2006b:37085)
- [PRL07] F. Przytycki and J. Rivera-Letelier, *Statistical properties of topological Collet-Eckmann maps*, Ann. Sci. École Norm. Sup. (4) **40** (2007), no. 1, 135–178. MR 2332354
- [PS75] W. Philipp and W. Stout, *Almost sure invariance principles for partial sums of weakly dependent random variables*, Mem. Amer. Math. Soc. 2 (1975), no. issue 2, 161, iv+140. MR 0433597
- [PUZ89] F. Przytycki, M. Urbański, and A. Zdunik, *Harmonic, Gibbs and Hausdorff measures on repellers for holomorphic maps. I*, Ann. of Math. (2) **130** (1989), no. 1, 1–40. MR 1005606
- [Sib99] N. Sibony, *Dynamique des applications rationnelles de $\mathbb{P}^k$* , Dynamique et géométrie complexes (Lyon, 1997), Panor. Synthèses, vol. 8, Soc. Math. France, Paris, 1999, pp. ix–x, xi–xii, 97–185. MR 1760844 (2001e:32026)
- [Tri78] H. Triebel, *Interpolation theory, function spaces, differential operators*, North-Holland Mathematical Library, vol. 18, North-Holland Publishing Co., Amsterdam, 1978. MR 503903 (80i:46032b)